



Experience effective Snow Mold control with CIVITAS TURF DEFENSE™

Set your turf up for a successful start of the Spring Season

For best results, make two 17 oz. applications 21 days apart. Applications should be made prior to turf dormancy. In most areas of the U.S., the second application should be complete by the middle to end of November. Based on historical pressure, tank mix one or more of your favorite Snow Mold control product(s) with the second application of CIVITAS TURF DEFENSE.

In the spring after the first mowing, start by making two to three 8.5 oz. applications on a 21 day interval for continued improved earlier spring playing conditions to your golfers. Tank mix with early season seedhead suppressants, insecticides and fungicides to promote a successful start to the season.



Salt Lake City, Utah - 1 x 17 oz. application in the fall.
Picture taken on April 7th 2015.



Early season applications result in continued dark green color from a plant that is functioning earlier in the season. This can often lead to the option of earlier season cultural practices.



CIVITAS TURF DEFENSE Ready-2-Mix 16 oz. + 1 oz. CIVITAS HARMONIZER Early & Late + 0.6 oz. Tebuconazole Late.



Tebuconazole 0.6 oz. Late.

MAXIMIZED TURF QUALITY - ENHANCED IPM EFFICIENCIES - IMPROVED DISEASE CONTROL



The active ingredient in CIVITAS TURF DEFENSE resembles the natural paraffins found on the surface of leaves and provides a physical barrier that helps prevent fungi from entering the turf plants. In addition, CIVITAS TURF DEFENSE features StrengthIN™ stress tolerance technology which increases the plants ability to tolerate both abiotic and biotic stresses.